

FORM PTO-1390 (Modified)
(REV 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES

211596US0PCT

DESIGNATED/ELECTED OFFICE (DO/EO/US)

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

CONCERNING A FILING UNDER 35 U.S.C. 371

09/926009

INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

PRIORITY DATE CLAIMED

PCT/EP00/01044

9 February 2000

12 February 1999

TITLE OF INVENTION

ARTIFICIAL PEPTIDES HAVING SURFACE ACTIVITY AND THE USE THEREOF IN THE PREPARATION OF ARTIFICIAL SURFACTANT

APPLICANT(S) FOR DO/EO/US

Tore CURSTEDT, et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☒ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☒ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☒ A copy of the International Search Report (PCT/ISA/210).

Items 13 to 20 below concern document(s) or information included:

13. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☐ Certificate of Mailing by Express Mail
23. ☒ Other items or information:

Amended Sheets (Pages 23, 24, 25, and Figure 1)

Notice of Priority

Request for Consideration of Documents Cited in the International Search Report

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

09/926009

INTERNATIONAL APPLICATION NO.

PCT/EP00/01044

ATTORNEY'S DOCKET NUMBER

211596US0PCT

24. The following fees are submitted..

CALCULATIONS PTO USE ONLY

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :

- ☐ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1000.00
- ☒ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00
- ☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

\$860.00

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☒ 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).

\$130.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	24 - 20 =	4	x \$18.00
Independent claims	1 - 3 =	0	x \$80.00

\$72.00

\$0.00

Multiple Dependent Claims (check if applicable). ☐

\$0.00

TOTAL OF ABOVE CALCULATIONS =

\$1,062.00

☐ Applicant claims small entity status. (See 37 CFR 1.27). The fees indicated above are reduced by 1/2.

\$0.00

SUBTOTAL =

\$1,062.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).

\$0.00

TOTAL NATIONAL FEE =

\$1,062.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). ☐

\$0.00

TOTAL FEES ENCLOSED =

\$1,062.00

Amount to be: refunded

\$

charged

\$

- a. ☒ A check in the amount of \$1,062.00 to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 15-0030 A duplicate copy of this sheet is enclosed.
- d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Telephone: (703)413-3000
Fax: (703)413-2220



22850

E. BEAUMONT
SIGNATURE REGISTRATION NUMBER 30,996

Norman F. Oblon

NAME

24,618

REGISTRATION NUMBER

DATE

August 13, 2001

09/926009

211596US-0 PCT

IN THE UNITED STATES PATENT & TRADEMARK OFFICE

IN RE APPLICATION OF: :

TORRE CURSTEDT ET AL : ATTN: APPLICATION DIVISION

SERIAL NO: NEW U.S. PCT APPLN :
(BASED ON PCT/EP00/01044)

FILED: HERewith :

FOR: ARTIFICIAL PEPTIDES HAVING
SURFACE ACTIVITY AND THE USE
THEREOF IN THE PREPARATION
OF ARTIFICIAL SURFACTANT

PRELIMINARY AMENDMENT

ASSISTANT COMMISSIONER FOR PATENTS
WASHINGTON, D.C. 20231

SIR:

Responsive to the Official Correspondence dated October 11, 2001, Applicants submit herewith a substitute Sequence Listing, a corresponding computer-readable Sequence Listing, and an amendment. Prior to examination on the merits, please amend the above-identified application as follows.

IN THE SPECIFICATION

Page 26 (Abstract), after the last line, beginning on the next page, please delete the original Sequence Listing attached at pages 1 to 3 and replace it with the substitute Sequence Listing attached hereto.

09926009 121101

Page 5, lines 23-25, please replace the paragraph with the following paragraph:

--As follows, according to a first aspect, the invention provides SP-C analogues having the following general formula (I) SEQ ID NO:1, using the one-letter amino acid code:--

Page 6, lines 21-26, please replace the paragraph with the following paragraph:

--Preferred peptides of Formula (I) have the following sequences:

(Ia) FGIPSSPVHLKRX₄BX₄BX₄BXGALLMGL (SEQ ID NO:2)

(Ib) FGIPSSPVHLKRX₅BX₅BX₄GALLMGL (SEQ ID NO:3)

(Ic) FGIPSSPVHLKRX₄BX₁₁GALLMGL (SEQ ID NO:4)

(Id) FGIPSSPVHLKRX₈BX₇GALLMGL (SEQ ID NO:5)

(Ie) FGIPSSPVHLKRX₁₁BX₄GALLMGL (SEQ ID NO:6)--

Page 7, lines 3-8, please replace the paragraph with the following paragraph:

--FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL [SP-C (LKS)] (SEQ ID NO:7)

FGIPSSPVHLKRLILLLIKLLILGALLMGL [SP-C (LKS)₁] (SEQ ID NO:8)

FGIPSSPVHLKRLILKLLLLILLILGALLMGL [SP-C (LKS)₂] (SEQ ID NO:9)

FGIPSSPVHLKRLILLLLLLLKLILLILGALLMGL [SP-C (LKS)₃] (SEQ ID NO:10)

FGIPSSPVHLKRLILLLLLLLIKLLILGALLMGL [SP-C (LKS)₄] (SEQ ID NO:11)

FGIPSSPVHLKRLILFLLLLFILLFLGALLMGL [SP-C (LFS)] (SEQ ID NO:12)-


Page 21, lines 27, to page 27, line 6, please replace the paragraph with the following paragraph:

--The sequence of human SP-C (SEQ ID NO: 13) is taken from Johansson, J., et al. (1988) FEBS Lett. **232**, 61-64 and that of SP-C(Leu) (SEQ ID NO: 14) from Nilsson, G., et al. (1999) Eur. J. Biochem, 255, 116-124). SP-C(LKS) (SEQ ID NO: 7) is based on the primary structure of SP-C but all Val residues at the positions 16-28 with the exception of position 17 are replaced with Leu residues, Lys residues have been introduced at positions 17,22, and 27, and the palmitoylated Cys at positions 5 and 6 are replaced with Ser.--

IN THE CLAIMS

Please amend the claims as follows:

--1. (Twice Amended) A SP-C analog having general formula (I) (SEQ ID NO:1), according to one-letter amino acid code:



wherein:

X is an amino acid selected from the group consisting of I, L, Nle (norleucine);

B is an amino acid selected from the group consisting of K, W, F, Y, Ornithine;

Z is S and can be optionally linked via ester or thio-ester bonds with acyl group containing 12-22 carbon atoms;

a is an integer from 1 to 19;

b is an integer from 1 to 19;

c is an integer from 1 to 21;

d is an integer from 0 to 20;

e is 0 or 1;

f is 0 or 1;

n is 0 or 1;

m is 0 or 1,

with the following conditions:

$n + m > 0$;

$f \geq e$;

$(X_aB)(X_bB)_n(X_cB)_mX_d$ is a sequence having a maximum of 22 amino acids.

2. (Amended) SP-C analogues according to claim 1, having formula (Ia) (SEQ ID NO:2)

(Ia) FGIPSSPVHLKRX₄BX₄BX₄BXGALLMGL

3. (Amended) SP-C analogues according to claim 1, having formula (Ib) (SEQ ID NO:3)

(Ib) FGIPSSPVHLKRX₅BX₅BX₄GALLMGL

4. (Amended) SP-C analogues according to claim 1, having formula (Ic) (SEQ ID NO:4)

(Ic) FGIPSSPVHLKRX₄BX₁₁GALLMGL

5. (Amended) SP-C analogues according to claim 1, having formula (Id) (SEQ ID NO:5)

(Id) FGIPSSPVHLKRX₈BX₇GALLMGL

6. (Amended) SP-C analogues according to claim 1, having formula (Ie) (SEQ ID NO:6)

(Ie) FGIPSSPVHLKRX₁₁BX₄GALLMGL

9. (Amended) SP-C analogues according to claim 8, selected from the group consisting of:

SP-C (LKS) FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL (SEQ ID NO:7)

SP-C (LKS)₁ FGIPSSPVHLKRLILKLLLLIKLLILGALLMGL (SEQ ID NO:8)

SP-C (LKS)₂ FGIPSSPVHLKRLILKLLLLLILLILGALLMGL (SEQ ID NO:9)

SP-C (LKS)₃ FGIPSSPVHLKRLILLLLLLKLILLILGALLMGL (SEQ ID NO:10)

SP-C (LKS)₄ FGIPSSPVHLKRLILLLLLLLIKLLILGALLMGL (SEQ ID NO:11)

SP-C (LFS) FGIPSSPVHLKRLILFLLLLFILLFLGALLMGL (SEQ ID NO:12)--

REMARKS

Claims 1-13 and 17-27 are active in the present application.

Applicants have now submitted a substitute Sequence Listing and a corresponding computer-readable Sequence Listing. Contents of the paper copy of the substitute Sequence Listing and the computer-readable Sequence Listing are identical. Support for all the sequences listed in the substitute Sequence Listing can be found in the present application. No new matter is introduced by the submission of the substitute Sequence Listing and the computer-readable Sequence Listing.

The specification and claims have been amended for clarity and to provide proper citation to the sequences in the substitute Sequence Listing. No matter is believed to be introduced by the amendments to the specification and claims.

In response to the Official Correspondence dated October 11, 2001, Applicants submit that the Declaration was properly filed on October 3, 2001 as evidenced by the copies of the Declaration and the date-stamped filing receipt submitted herewith.

Applicants submit that this application is in condition for allowance. Early notice to this effect is earnestly solicited.

Respectfully submitted,

OBLON, SPIVAK, McCLELLAND,
MAIER & NEUSTADT, P.C.



Norman F. Oblon
Attorney of Record
Registration No. 24,618

Daniel J. Pereira, Ph.D.
Registration No. 45,518

J. Derek Mason, Ph.D.
Registration No. 35,270



22850

(703) 413-3000
Fax #: (703) 413-2220
I:\atty\Twb\211596US-PR.wpd

Marked-Up Copy

Serial No: 09/926,009

Amendment Filed on:

December 11, 2001

IN THE SPECIFICATION

Page 5, lines 23-25, please replace the paragraph with the following paragraph:

--As follows, according to a first aspect, the invention provides SP-C analogues having the following general formula (I) SEQ ID NO:1, using the one-letter amino acid code:--

Page 6, lines 21-26, please replace the paragraph with the following paragraph:

--Preferred peptides of Formula (I) have the following sequences:

(Ia) FGIPSSPVHLKRX₄BX₄BX₄BXGALLMGL (SEQ ID NO:2)

(Ib) FGIPSSPVHLKRX₅BX₅BX₄GALLMGL (SEQ ID NO:3)

(Ic) FGIPSSPVHLKRX₄BX₁₁GALLMGL (SEQ ID NO:4)

(Id) FGIPSSPVHLKRX₈BX₇GALLMGL (SEQ ID NO:5)

(Ie) FGIPSSPVHLKRX₁₁BX₄GALLMGL (SEQ ID NO:6)--

Page 7, lines 3-8, please replace the paragraph with the following paragraph:

--FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL [SP-C (LKS)] (SEQ ID NO:7)

FGIPSSPVHLKRLILLLIKLLILGALLMGL [SP-C (LKS)₁] (SEQ ID NO:8)

FGIPSSPVHLKRLILKLLLLLILLILGALLMGL [SP-C (LKS)₂] (SEQ ID NO:9)

FGIPSSPVHLKRLILLLLLLLKLILLILGALLMGL [SP-C (LKS)₃] (SEQ ID NO:10)

FGIPSSPVHLKRLILLLLLLLIKLLILGALLMGL [SP-C (LKS)₄] (SEQ ID NO:11)


FGIPSSPVHLKRLILFLLLFILLFLGALLMGL [SP-C (LFS)] (SEQ ID NO:12)-

Page 21, lines 27, to page 27, line 6, please replace the paragraph with the following paragraph:

--The sequence of human SP-C (SEQ ID NO: 13) is taken from Johansson, J., et al. (1988) FEBS Lett. **232**, 61-64 and that of SP-C(Leu) (SEQ ID NO: 14) from Nilsson, G., et al. (1999) Eur. J. Biochem, 255, 116-124). SP-C(LKS) (SEQ ID NO: 7) is based on the primary structure of SP-C but all Val residues at the positions 16-28 with the exception of position 17 are replaced with Leu residues, Lys residues have been introduced at positions 17,22, and 27, and the palmitoylated Cys at positions 5 and 6 are replaced with Ser.--

IN THE CLAIMS

--1. (Twice Amended) A SP-C analog having general formula (I) (SEQ ID NO:1), according to one-letter amino acid code:



wherein:

X is an amino acid selected from the group consisting of I, L, Nle (norleucine);

B is an amino acid selected from the group consisting of K, W, F, Y, Ornithine;

Z is S and can be optionally linked via ester or thio-ester bonds with acyl group containing 12-22 carbon atoms;

a is an integer from 1 to 19;

b is an integer from 1 to 19;

c is an integer from 1 to 21;

d is an integer from 0 to 20;

e is 0 or 1;

f is 0 or 1;

n is 0 or 1;

m is 0 or 1,

with the following conditions:

$n + m > 0$;

$f \geq e$;

$(X_aB)(X_bB)_n(X_cB)_mX_d$ is a sequence having a maximum of 22 amino acids.

2. (Amended) SP-C analogues according to claim 1, having formula (Ia) (SEQ ID NO:2)

(Ia) FGIPSSPVHLKRX₄BX₄BX₄BXGALLMGL

3. (Amended) SP-C analogues according to claim 1, having formula (Ib) (SEQ ID NO:3)

(Ib) FGIPSSPVHLKRX₅BX₅BX₄GALLMGL

4. (Amended) SP-C analogues according to claim 1, having formula (Ic) (SEQ ID NO:4)

(Ic) FGIPSSPVHLKRX₄BX₁₁GALLMGL

5. (Amended) SP-C analogues according to claim 1, having formula (Id) (SEQ ID NO:5)

(Id) FGIPSSPVHLKRX₈BX₇GALLMGL

6. (Amended) SP-C analogues according to claim 1, having formula (Ie) (SEQ ID NO:6)

(Ie) FGIPSSPVHLKRX₁₁BX₄GALLMGL

9. (Amended) SP-C analogues according to claim 8, selected from the group consisting of:

SP-C (LKS) FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL (SEQ ID NO:7)

SP-C (LKS)₁ FGIPSSPVHLKRLILKLLLLIKLLILGALLMGL (SEQ ID NO:8)

SP-C (LKS)₂ FGIPSSPVHLKRLILKLLLLILLLILGALLMGL (SEQ ID NO:9)

SP-C (LKS)₃ FGIPSSPVHLKRLILLLLLLKLILLILGALLMGL (SEQ ID NO:10)

SP-C (LKS)₄ FGIPSSPVHLKRLILLLLLLLIKLLILGALLMGL (SEQ ID NO:11)

SP-C (LFS) FGIPSSPVHLKRLILFLLLFILLFLGALLMGL (SEQ ID NO:12)--

211596US-0PCT

IN THE UNITED STATES PATENT & TRADEMARK OFFICE

IN RE APPLICATION OF :
TORE CURSTEDT ET AL : ATTN: APPLICATION DIVISION
SERIAL NO: NEW U.S. PCT APPLN :
(Based on PCT/EP00/01044)
FILED: HEREWITH :
FOR: ARTIFICIAL PEPTIDES HAVING :
SURFACE ACTIVITY AND THE
USE THEREOF IN THE
PREPARATION OF ARTIFICIAL
SURFACTANT

PRELIMINARY AMENDMENT

ASSISTANT COMMISSIONER FOR PATENTS
WASHINGTON, D.C. 20231

SIR:

Prior to examination on the merits, please amend the above-identified application as follows:

IN THE CLAIMS

Please delete Claims 14-16.

Please amend the claims as shown on the marked-up copy following this amendment to read as follows:

1. (Amended) A SP-C analog having general formula (I), according to one-letter amino acid code:



wherein:

X is an amino acid selected from the group consisting of I, L, Nle (norleucine);

B is an amino acid selected from the group consisting of K, W, F, Y, Ornithine;

Z is S and can be optionally linked via ester or thio-ester bonds with acyl group containing 12-22 carbon atoms;

a is an integer from 1 to 19;

b is an integer from 1 to 19;

c is an integer from 1 to 21;

d is an integer from 0 to 20;

e is 0 or 1;

f is 0 or 1;

n is 0 or 1;

m is 0 or 1,

with the following conditions:

$n + m > 0$;

$f \geq e$;

$(X_a B)(X_b B)_n (X_c B)_m X_d$ is a sequence having a maximum of 22 amino acids.

7. (Amended) A SP-C analog according to Claim 1, in which Ser residues are acylated.

8. (Amended) A SP-C analog according to Claim 1, in which B is Lysine or Phenylalanine and X is Leucine, Isoleucine or Norleucine.

10. (Amended) A synthetic surfactant comprising at least one SP-C analogue as claimed in Claim 1 in admixture with lipids and phospholipids.

12. (Amended) A synthetic surfactant according to Claim 10, further comprising SP-B or an active derivative thereof or a polymyxin.

13. (Amended) A synthetic surfactant according to Claim 10, in form of solution, dispersion, suspension, dry powder.

Please add new Claims 17-27 as follows:

17. (New) The SP-C analogue as claimed in claim 1 wherein the $(X_aB)(X_bB)_n(X_cB)_mX_d$ sequence has from 10 to 22 amino acids.

18. (New) The SP-C analogue as claimed in claim 7 wherein the Ser residues are acylated with palmitoyl groups.

19. (New) A pharmaceutically active synthetic surfactant comprising the SP-C analogues of Claim 1.

20. (New) A method of treating a surfactant deficiency comprising administering the SP-C analogues of Claim 1.

21. (New) A pharmaceutically active synthetic surfactant comprising the surfactant of Claim 10, wherein said surfactant comprises polymyxin.

22. (New) A pharmaceutically active synthetic surfactant comprising the surfactant of Claim 10, wherein said surfactant comprises polymyxin B.

23. (New) A method of treating surfactant deficiencies or dysfunction, or serious otitis media comprising administering the surfactant of Claim 10 wherein said surfactant comprises polymyxin.

24. (New) A method of treating surfactant deficiencies or dysfunction, or serious otitis media comprising administering the surfactant of Claim 10 wherein said surfactant comprises polymyxin B.

25. (New) The method of treating a surface deficiency of Claim 20, wherein said surfactant deficiency is respiratory distress syndrome.


26. (New) The method of treating a surface deficiency of Claim 23, wherein said surfactant deficiency is respiratory distress syndrome.

27. (New) The method of treating a surface deficiency of Claim 24, wherein said surfactant deficiency is respiratory distress syndrome.

REMARKS

Claims 1-13 and 17-27 are active in the present application. Claims 1-13 were amended to remove multiple dependencies. Claims 14-16 were canceled. New Claims 17-27 replace canceled Claims 14-17. Support for new claims is found in the original claims and in the specification on page 1, lines 5-12 and the examples. No new matter is believed to have been added. An action on the merits and allowance of claims is solicited.

Respectfully submitted,
OBLON, SPIVAK, McCLELLAND,
MAIER & NEUSTADT, P.C.



Norman F. Oblon
Attorney of Record
Registration No. 24,618

Daniel J. Pereira, Ph.D.
Registration No. 45,518



22850

(703) 413-3000
Fax #: (703) 413-2220
DJPER/kst
I:\atty\SUKOS\211596us-pr.wpd

Marked-Up Copy


Serial No:

Amendment Filed on:

08-13-01

IN THE CLAIMS

--1. (Amended) A SP-C analog [SP-C analogues] having general formula (I),
according to one-letter
amino acid code:



wherein:

X is an amino acid selected from the group consisting of I, L, Nle (norleucine);

B is an amino acid selected from the group consisting of K, W, F, Y, Ornithine;

Z is S and can be optionally linked via ester or thio-ester bonds with acyl group
containing 12-22 carbon atoms;

a is an integer from 1 to 19;

b is an integer from 1 to 19;

c is an integer from 1 to 21;

d is an integer from 0 to 20;

e is 0 or 1;

f is 0 or 1;

n is 0 or 1;

m is 0 or 1,

with the following conditions:

$$n + m > 0;$$

$$f \geq e;$$

$(X_aB)(X_bB)_n(X_cB)_mX_d$ is a sequence having a maximum of 22 amino acids[, preferably from 10 to 22 amino acids].

7. (Amended) A SP-C analog [SP-C analogues] according to [claims 1-6] Claim 1, in which Ser residues are acylated[preferably with palmitoyl groups].

8. (Amended) A SP-C analog [SP-C analogues] according to [claims 1-7] Claim 1, in which B is Lysine or Phenylalanine and X is Leucine, Isoleucine or Norleucine.

10. (Amended) A synthetic surfactant comprising at least one SP-C analogue [of formula (I)] as claimed in Claim 1 in admixture with lipids and phospholipids.

12. (Amended) A synthetic surfactant according to [claims 10-11] Claim 10, further comprising SP-B or an active derivative thereof or a polymyxin.

13. (Amended) A synthetic surfactant according to [claims 10-12] Claim 10, in form of solution, dispersion, suspension, dry powder.--

Claims 14-16 (Canceled).

Claims 17-27 (New).

**ARTIFICIAL PEPTIDES HAVING SURFACE ACTIVITY AND THE
USE THEREOF IN THE PREPARATION OF ARTIFICIAL
SURFACTANT**

The present invention provides new artificial peptides having surface activity. In particular, the invention provides SP-C analogues which, once combined with suitable lipids, are particularly effective in reducing surface tension at the air-liquid interface.

5 Thus, the peptides of the invention may be used in combination with lipids, and optionally in combination with SP-B or an active analogue thereof or a substitute of SP-B, for preparing artificial surfactants useful in the treatment of respiratory distress syndrome (RDS), other surfactant deficiencies or dysfunction,
10 related pulmonary diseases such as pneumonia, bronchitis, asthma, meconium aspiration syndrome and also other diseases such as serous otitis media (glue ear).

Background of the invention

Pulmonary surfactant reduces surface tension at the air-liquid
15 interface of the alveolar lining, preventing the lungs from collapsing at end expiration. Surfactant deficiency is a common disorder in premature infants and causes respiratory distress syndrome (RDS), which can be effectively treated with natural surfactants extracted from animal lungs (Fujiwara, T. and
20 Robertson B. (1992) *In*: Robertson, B., van Golde, L.M.G. and Batenburg, B. (eds) Pulmonary Surfactant: From Molecular Biology to Clinical Practice Amsterdam, Elsevier, pp. 561-592). The main constituents of these surfactant preparations are phospholipids such as 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC),

phosphatidylglycerol (PG) and the hydrophobic surfactant proteins B and C (SP-B and SP-C). The hydrophilic surfactant proteins SP-A and SP-D which are C-type (Ca^{2+} -dependent) collagenous lectins and thought to act primarily in the host-defence system, are normally not included in the surfactant preparations due to the organic solvent extraction procedures employed.

SP-B and SP-C constitute only about 1-2 % of the surfactant mass, but are still able to exercise dramatic improvements on surface activity, compared to pure lipid preparations (Curstedt, T. et al. (1987) *Eur. J. Biochem.* **168**, 255-262; Takahashi, A., Nemoto, T. and Fujiwara, T. (1994) *Acta Paediatr. Jap.* **36**, 613-618). The primary and secondary structures of SP-B and SP-C and a tertiary structure of SP-C in solution have been determined (see 4). SP-B is composed of two identical polypeptide chains of 79 amino acids, connected with an interchain disulphide bridge (Curstedt, T. et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**, 2985-2989; Johansson, J., Curstedt, T. and Jörnvall, H. (1991) *Biochemistry* **30**, 6917-6921). Each monomeric chain has three intrachain disulphide bridges and at least four amphipathic helices exhibiting one polar and one unpolar face through which SP-B may interact with two lipid bilayers and bring them into close proximity (Andersson, M. et al. (1995) *FEBS Lett.* **362**, 328-332). SP-C is a lipoprotein composed of 35 amino acid residues with an α -helical domain between residues 9-34 (Johansson, J. et al. (1994) *Biochemistry* **33**, 6015-6023). The helix is composed mostly of valyl-residues and is embedded in a lipid bilayer and oriented in parallel with the lipid acyl chains (Vandenbussche, et al. (1992)

Eur. J. Biochem. **203**, 201-209). Two palmitoyl groups are covalently linked to cysteine residues in positions 5 and 6 in the N-terminal part of the peptide (Curstedt, T. et al. (1990) Proc. Natl. Acad. Sci. U.S.A. **87**, 2985-2989). The two conserved positively charged residues, arginine and lysine, at positions 11 and 12, possibly interact with the negatively charged head groups of the lipid membrane, thus increasing its rigidity. The rigidity of the lipid-peptide interaction may be decreased towards the C-terminal end, since it contains small or hydrophobic residues only, making this part potentially more mobile in a phospholipid bilayer. SP-C is thought to influence the thickness and fluidity of the surrounding lipids *via* the extremely stable poly-valyl helix (Johansson, J. and Curstedt, T. (1997) Eur. J. Biochem. **244**, 675-693)

State of the art

Since surfactant preparations obtained from animal tissue present some drawbacks, like their availability in limited amounts and the possibilities that they contain infectious agents and induce immunological reactions, attempts have been made to create artificial surfactants (Johansson, J. and Curstedt, T. (1997) Eur. J. Biochem. **244**, 675-693; Johansson, J. et al. (1996) Acta Paediatr. **85**, 642-646), usually from synthetic lipids and hydrophobic proteins.

Previous work has demonstrated that synthetic SP-C may not fold like the native peptide into an α -helical conformation necessary for optimal surface activity (Johansson, J. et al. (1995) Biochem. J. **307**, 535-541), and therefore do not interact properly with the surfactant lipids.

Consequently, synthetic SP-C analogues do not fold like the

native peptide and do not interact properly with the surfactant lipids. To circumvent this problem, several attempts have been made to modify the sequence, for instance by replacing all helical Val residues in native SP-C with Leu, which strongly favour α -helical conformation. The corresponding transmembranous analogue, SP-C(Leu) showed good spreading at an air-liquid interface when combined with DPPC:PG:PA (68:22:9) (w/w/w). However, the maximum surface tension value during cyclic surface compression (γ_{\max}) was significantly higher than that of native surfactant. It was furthermore not possible to prepare lipid-peptide mixtures of higher concentrations than about 20 mg/ml, probably due to formation of peptide oligomers (Nilsson, G. et al. (1998) Eur. J. Biochem. 255, 116-124). Others have synthesised bioactive polyleucine SP-C analogues of different lengths (Takei, T. et al. (1996) Biol. Pharm. Bull. **19**, 1550-1555). In the latter studies neither self oligomerisation nor problems in producing samples of high lipid concentration were reported.

Different publications deal with the problem of providing peptidic analogues of natural surfactant peptides, giving a number of different solutions. Among those publications, WO93 21225, EP 733 645, WO96 17872, in the name of Tokyo Tanabe, disclose peptides analogues of natural SP-C, which in general differ from the native peptide regarding the sequence of the N-terminal part.

Scripps Research Institute patent applications WO89 06657 and WO92 22315 disclose SP-B analogues having alternating hydrophobic and hydrophilic amino acid residues. Among others, a peptide alternating Leucine and lysine residues (KL₄) is claimed.

Clercx A. et al., Eur. J. Biochem 229, 465-72, 1995, disclose

peptides of different lengths corresponding to the N-terminal of porcine SP-C and hybrid peptides derived from porcine SP-C and bacteriorhodopsin.


Johansson J. et al., Biochem. J. 307, 535:41, 1995, disclose synthetic peptides that differ from the native porcine SP-C by the substitution of some amino acids.

WO89/ 04326 in the name of California Biotechnology - Byk Gulden, and WO91/18015 in the name of California Biotechnology - Scios Nova, disclose SP-C analogues containing an initial N-terminal sequence in which the two Cys of natural SP-C are replaced by two Ser.

Description of the invention

It has now been found that SP-C analogue peptides which combine the following features: i) substitution of Val residues with other neutral and hydrophobic residues; ii) substitution of Cys residues with Ser residues; iii) replacement of some of the neutral amino acid residues with bulky or polar residues, show particularly favourable properties for surface tension reduction. In particular it has been found that the latter feature, in virtue of the positive charges conferred by the polar residues or the steric hindrance conferred by the bulky substituents, allow to avoid self-oligomerisation.

As follows, according to a first aspect, the invention provides SP-C analogues having the following general formula (I), using the one-letter amino acid code:



wherein:

X is an amino acid selected from the group consisting of

V, I, L, Nle (norleucine);

B is an amino acid selected from the group consisting of Ornithine, K, I, W, F, Y, Q, N;

5 Z is an amino acid selected from the group consisting of S, C, F where Ser or Cys residues are optionally linked via ester or thio-ester bonds with acyl groups containing 12-22 carbon atoms linked.

a is an integer from 1 to 19

b is an integer from 1 to 19

10 c is an integer from 1 to 21

d is an integer from 0 to 20

e is 0 or 1

f is 0 or 1

n is 0 or 1

15 m is 0 or 1

with the conditions:

- $n + m > 0$,

- $f \geq e$;

20 - $(X_a B)_n (X_b B)_n (X_c B)_m X_d$ is a sequence having a maximum of 22 amino acids, preferably from 10 to 22.

Preferred peptides of Formula (I) have the following sequences:

(Ia) FGIPSSPVHLKRX₄BX₄BX₄BXGALLMGL

(Ib) FGIPSSPVHLKRX₅BX₅BX₄GALLMGL

(Ic) FGIPSSPVHLKRX₄BX₁₁GALLMGL

25 (Id) FGIPSSPVHLKRX₈BX₇GALLMGL

(Ie) FGIPSSPVHLKRX₁₁BX₄GALLMGL

Among the sequences (Ia) - (Ie), those having B = Lys or Phe and X = Leu, Ile or Nle are preferred.

According to preferred embodiments, peptides of formula (Ia) -
(If) have the following sequences, respectively:

FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL [SP-C (LKS)]

FGIPSSPVHLKRLILLLKLLLLIKLLILGALLMGL [SP-C (LKS)₁]

5 FGIPSSPVHLKRLILKLLLLLLILGALLMGL [SP-C (LKS)₂]

FGIPSSPVHLKRLILLLLLLKLLLLILGALLMGL [SP-C (LKS)₃]

FGIPSSPVHLKRLILLLLLLLIKLLILGALLMGL [SP-C (LKS)₄]

FGIPSSPVHLKRLILFLLLLFILLFLGALLMGL [SP-C (LFS)]

10 In a more preferred embodiment of the invention, the Ser
residues are covalently linked with acyl groups containing 12-22
carbon atoms.

Peptides of formula (I) may be prepared by synthetic methods
or recombinant techniques.

15 Conventional synthetic methods are described, for instance,
in Schroeder et al., "The peptides", vol. 1, Academic Press, 1965;
Bodanszky et al., "Peptide synthesis", Interscience Publisher,
1996; Baramy & Merrifield, "The peptides; Analysis, Synthesis,
Biology", vol. 2, chapter 1, Academic Press, 1980. Said techniques
include peptide synthesis in solid phase, in solution, organic
20 chemistry synthetic methods, or any combination thereof.

S- or O- acylated peptides are preferably synthesized by
treatment of the non-acylated peptides with acyl chloride in neat
trifluoroacetic acid as described in Yousefi-Salakdeh et al.
Biochem J 1999, 343, 557-562. After synthesis and purification,
25 the synthetic peptides were biochemically and biophysically
characterised, as reported in the following section "Examples".

The activity of the peptides of the invention in reducing
surface tension has been evaluated in combination with lipids and

phospholipids, SP-B, analogues of SP-B or substitutes of SP-B. In particular, the peptides have been combined with DPPC (1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine)/PG (phosphatidylglycerol)/PA (palmitic acid) with or without SP-B, an active analogue thereof and polymyxins.

The results of pulsating-bubble surface activity tests clearly show that the synthetic peptides according to the present invention strongly decrease minimum and maximum surface tension during cyclic surface compression (γ_{\min} and γ_{\max}) to values comparable with those obtained using surfactants from natural sources.

The addition of SP-B or active analogue thereof to the mixture peptide/lipids-phospholipids gave particularly favourable results. Furthermore it has been surprisingly found that polymyxins, in particular polymyxin B, act as substituents of SP-B and their addition gave comparable results to those achieved with SP-B.

According to a second aspect, the invention provides a synthetic surfactant comprising one or more peptides of formula (I), in admixture with lipids and/or phospholipids and optionally SP-B, an active derivative thereof or polymyxins. Suitable lipids/phospholipids may be selected from the group consisting of phosphatidylcholines (preferably DPPC) PG, PA, triacylglycerols, sphingomyelin.

In an even more preferred embodiment of the invention, surfactant mixtures containing the peptide in which palmitoyl chains are O-covalently linked to the Ser residues should be used. It has been found that surfactant mixtures containing a dipalmitoylated form of the reference peptide (SP-C(Leu)) exhibit

higher surface film stability and increased size of the surface-associated lipid reservoir, compared to mixtures containing the corresponding non-palmitoylated peptide, as measured by captive bubble system. In the samples containing 5% of dipalmitoylated peptide, the γ_{\min} was below 1.5 mN/m and the films very stable, as the surface tension increased by less than 0.5 mN/m within 10 min at constant bubble volume. On the contrary, the γ_{\min} for the non-palmitoylated peptide was approx. 5 mN/m and the films were less stable as observed by frequent bubble clicking at low surface tensions. Moreover, after subphase depletion for samples that contain non-palmitoylated peptide, the ability to reach near zero stable surface tension was lost after a few adsorption steps, whereas with the dipalmitoylated peptide the film quality did not deteriorate even after more than 10 expansion steps and the incorporation of reservoir material equivalent to more than two monolayers. The improved surface activity of dipalmitoylated peptides was also demonstrated by pulsating bubble surfactometer. In addition, the presence of acyl groups was found to further reduce the tendency to form oligomers. This finding is very important, as during preparation of artificial surfactants, peptide oligomerisation has been found to hinder preparation of the mixtures at higher concentrations than 20 mg/ml (Nilsson *et al.* Eur J Biochem 1998, 255, 116-124).

The synthetic surfactant may be prepared by mixing solutions or suspensions of peptides and lipids and by subsequently drying the mixture.

At the occurrence, the dry mixture may be suspended, dispersed or administered as such to subjects in need of treatment

for surfactant deficiency.

The synthetic surfactant will be preferably administered endotracheally or via aerosol. The latter form of administration will require the combination of small particles of surfactant with
5 suitable inert propellant. Other forms of administration, like nebulization or spraying of stable solutions/suspensions of surfactant are also included within the scope of the invention.

According to a further aspect, the invention provides the use of the described peptides for the preparation of a surfactant agent
10 to be used in all cases of adult or neonatal surfactant deficiency or dysfunction, related pulmonary diseases such as pneumonia, bronchitis, asthma, meconium aspiration syndrome and also other diseases such as serous otitis media (glue ear).

Typically, the surfactant agent will be used, preferably upon
15 endotracheal administration, in the treatment of respiratory distress syndrome which frequently affects premature infants.

The following examples illustrate the invention in more details.

Example 1

20 **Peptide synthesis and purification**

An analogue of SP-C, SP-C(LKS) (Fig. 1) was synthesised by use of stepwise solid phase technology and the *tert*-butyloxycarbonyl chemistry (Kent, S.B.H. (1988) Annu. Rev. Biochem. **57**, 957-989) in an Applied Biosystems 430A instrument.
25 Cleavage of the resin-peptide bond and deprotection of the side-chains were carried out in anhydrous hydrogen fluoride/methoxybenzene/dimethylsulfide, 10:1:1 (v/v/v) for 1.5 h at 0°C. Protecting groups and scavengers were removed by

repeated extraction with diethyl ether and the peptide was subsequently extracted from the resin by dichloromethane/trifluoroacetic acid (TFA) 3:1 (v/v) followed by rotary evaporation. The crude peptide extract was redissolved at a concentration of 100 mg/ml in chloroform/methanol 1:1 (v/v) containing 5% H₂O. An aliquot of 10 mg was applied on a Sephadex LH-60 column (40 x 1 cm) in the same solvent (Curstedt, T. et al. (1987) Eur. J. Biochem. **168**, 255-262). Fractions of 2.5 ml were collected and absorbencies at 214 and 280 nm were measured. Identification and quantitation were performed by amino acid analysis.

For acylation, the purified peptide (typically about 5 mg) is dried, dissolved in distilled TFA (100 µl) and acyl chloride (10-20 equivalents compared to peptide) is added. After 10 minutes the reaction is quenched with 80% aqueous ethanol (1.9 ml). Purification of acyl peptides is performed using chromatography over Lipidex 5000 in ethylene chloride/methanol 1:4 (v/v) followed by reversed-phase HPLC over a C18 column using a linear gradient of 2-propanol/0.1% TFA running into 60% (aqueous) methanol/0.1% TFA or 75% (aqueous) ethanol/0.1% TFA.

Example 2

Biochemical characterisation

The purity of the peptide was checked by sodium-dodecylsulphate (SDS) polyacrylamide gel electrophoresis (PAGE) (Phast-system, Pharmacia, Sweden) and by reversed phase high performance liquid chromatography (HPLC), using a C₁₈ column and a linear gradient of 60 % aqueous methanol /0.1 % TFA and isopropanol/ 0.1 % TFA (Gustafsson, M. et al. (1997) Biochem. J.

326, 799-806).

Molecular masses were determined by matrix-assisted laser desorption ionisation-time-of-flight (MALDI-TOF) mass spectrometry (Lasermat 2000, Finnigan MAT) calibrated with vasoactive intestinal peptide (M_r 3326.8).

Peptide secondary structure was investigated using circular dichroism (CD) spectroscopy (Jasco-720 Jasco, Japan). After solubilisation with trifluoroethanol (TFE) spectra were recorded from 260 to 184 nm with a scan speed of 20 nm/min and a resolution of 2 data points/nm. The residual molar ellipticity was calculated and expressed in $\text{kdeg} \times \text{cm}^2/\text{dmol}$. Molar ellipticities at 208 and 222 nm were utilised for estimating the content of helical structure (Barrow, C.J. et al. (1992) J. Mol. Biol. **225**, 1075-1093).

Secondary structure investigations of SP-C(LKS) using CD spectroscopy showed a spectrum typical for α -helical peptides and an α -helical content of approximately 75 % was estimated from the 208 nm and 222 nm minima. The secondary structure remained stable following stepwise dilution with H_2O until 12 % TFE provided that the peptide was solubilized in neat TFE.

SDS-PAGE of SP-C(LKS) showed a single band similar to native SP-C while SP-C(Leu) which lacks Lys in the helical part forms oligomers. In contrast to our experience with SP-C(Leu)/lipid mixtures, which are difficult to solubilize in higher concentrations than 20 mg/ml (Nilsson, G. et al. (1998) Eur. J. Biochem. **255**, 116-124.), it was possible to make a SP-C(LKS)/lipid mixture with a lipid concentration of 80 mg/ml and a polypeptide/lipid ratio of 0.03.

Example 3

Preparation of peptide/lipid mixtures

DPPC, PG and PA were all purchased from Sigma Chemical Co. (St Louis, MO). The lipids, dissolved in chloroform/methanol 98:2 (v/v), were mixed in the proportions DPPC:PG:PA 68:22:9 (w/w/w) or DPPC/PG 7:3 (w/w). Surfactant preparations were prepared by adding SP-C(LKS) alone or SP-C(LKS) and SP-B, to each of the lipid mixtures, at total polypeptide/lipid weight ratios of 0-0.05. The mixtures were evaporated under nitrogen and resuspended in 150 mmol/l NaCl or in 10 mmol/l Hepes buffer pH 6.9 containing 140 mmol/l NaCl and 2.0 mmol/l CaCl_2 , at lipid concentrations of 10 — 80 mg/ml. Repeated freezing and sonication (50 W, 48 kHz) were performed until homogeneous suspensions were achieved. In some cases the final suspensions were incubated at 45°C for 1 h.

Surfactant preparations suspended in 150 mmol/l NaCl have a pH of 3.5-5.5. The lower pH-values 3.5-4.5 were observed in preparations containing SP-B. Since native SP-B is purified using acidified organic solvents (Curstedt, T. et al. (1987) Eur. J. Biochem. **168**, 255-262) small amounts of acid may remain in the preparations. Near physiological pH was obtained by suspending the surfactant preparation in Hepes buffer pH 6.9, containing 140 mmol/l NaCl and 2 mmol/l CaCl_2 (Table 1). Compared to the corresponding preparations in unbuffered saline there were no changes of γ_{max} or γ_{min} when DPPC/PG 7:3 (w/w) was used as the lipid mixture. However when PA was included in the lipid mixture both γ_{max} and γ_{min} increased at the higher pH (Tables 1 and 2).

Table 1: Surface properties of preparations of artificial surfactant in physiological saline solution

Measurements were carried out directly after the preparation of the samples or after incubation for 1 hour at 45° C. The phospholipids conc. was 10 mg/ml in NaCl 150 mmol/l. The recordings were obtained at different periods of time with a pulsating bubble surfactometer at 37° C, 50% surface compression and at a rate of 40 cycles per min. The values are the mean (standard deviation) of 3-5 measurements. Abbreviations are defined in the text.

Surfactant preparation			Surface tension (mN/m)						
SP-C (LKS) (% w/w)	SP-B (% w/w)	Phospholipids	Incubation temperature	7,5 s		1 min		5 min	
				γ_{min}	γ_{max}	γ_{min}	γ_{max}	γ_{min}	γ_{max}
3	-	DPPC/PG/PA	-	<1	41(1)	<1	41(1)	<1	41(0)
3	-	DPPC/PG/PA	45°C	<1	41(1)	<1	41(1)	<1	41(1)
3	2	DPPC/PG/PA	-	<1	33(2)	<1	33(2)	<1	33(2)
3	2	DPPC/PG/PA	45°C	<1	34(1)	<1	34(2)	<1	25(2)
3	-	DPPC/PG	-	12	39(5)	14(4)	42(4)	9(5)	42(2)
3	-	DPPC/PG	45°C	8(3)	35(5)	9(3)	39(5)	6(4)	42(3)
3	2	DPPC/PG	-	2(1)	31(1)	2(1)	31(3)	2(1)	33(1)
3	2	DPPC/PG	45°C	3(3)	29(3)	1(1)	33(2)	1(0)	36(1)
3	1	DPPC/PG	-	<1	24(4)	1(2)	26(4)	<1	31(1)
3	0.5	DPPC/PG	45°C	4(2)	29(1)	4(3)	29(2)	3(1)	34(2)

Table 2: Surface properties of preparations of artificial surfactant in buffered salt solution

Measurements were carried out on samples containing phospholipids at the conc. of 10 mg/ml in Hepes buffer (pH 6.9), in turn containing NaCl 140 mmol/l and CaCl₂ 2.0 mmol/l. The recordings were obtained at different periods of time with a pulsating bubble surfactometer at 37° C, 50% surface compression and at a rate of 40 cycles per min. The values are the mean (standard deviation) of 3-5 measurements. Abbreviations are defined in the text.

Surfactant preparation		Surface tension (mN/m)			
SP-C (LKS) (% w/w)	SP-B (% w/w)	Phospholipids	7,5 s ----- γ_{\min} γ_{\max}	1 min ----- γ_{\min} γ_{\max}	5 min ----- γ_{\min} γ_{\max}
3	-	DPPC/PG/PA	4(1) 44(2)	5(2) 47(2)	7(2) 50(1)
3	2	DPPC/PG/PA	3(3) 38(3)	4(4) 40(2)	3(3) 44(2)
3	-	DPPC/PG	15(3)	16(2) 42(3)	13(3) 44(3)
3	2	DPPC/PG	39(4) 26(3)	1(1) 29(3)	<1 35(1)

Example 4.

Preparation of phospholipid mixtures with SP-C(LKS) and polymyxin B.

DPPC and PG were purchased from Sigma Chemical Co (St Louis, MO). The phospholipids, dissolved in chloroform/methanol 98:2 (v/v), were mixed in the proportions DPPC/PG 7:3 (w/w). SP-C (LKS) was added to the phospholipid mixtures, at a total polypeptide/phospholipid weight ratio of 0.03. The mixtures were evaporated under nitrogen and resuspended at room temperature in 10 mmol/l Hepes buffer pH 6,9 containing 140 mmol/l NaCl and 2.0 mmol/l CaCl₂ or in the same buffer containing 0.01% polymyxin B (PxB) (Sigma Chemical Co, St Louis, MO). Repeated freezing and sonication (50 W, 48 kHz) were performed until homogeneous suspensions were achieved. The final phospholipid concentration for both preparations was 10 mg/ml. Addition of PxB decreased both γ_{min} and γ_{max} and optimal surface activity was obtained (Table 3).

Table 3. Surface properties of artificial surfactant with and without polymyxin B

The recordings were obtained at different periods of time with a pulsating bubble surfactometer at 37°C and 50% surface compression at a rate of 40 cycles per min. The values are the mean (SD) of 5-11 measurements. Abbreviations are defined in the text.

Surfactant preparation			Surface tension (mN/m)					
SP-C (LKS) (% w/w)	PxB (% w/w)	Phospholipids	7,5 s		1 min		5 min	
			γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}
3 3	- 1	DPPC/PG DPPC/PG	15(3) 3(2)	39(4) 29(3)	16(2) 2(2)	42(3) 31(4)	13(3) 1(1)	44(3) 34(2)

Example 5

Biophysical characterisation

Surface spreading kinetics were measured at about 34-37 °C with a Wilhelmy surface balance (Biegler, Vienna, Austria). Surface tension was monitored for 10 min using a platinum plate connected to a strain gauge and inserted 1 mm in to a hypophase of 20 ml of 150 mmol/l NaCl in a teflon trough. The suspensions were added as droplets, totally 1 mg of lipids, onto the hypophase, 4 cm from the platinum plate.

Kinetic measurements of 3 weight % SP-C(LKS) in DPPC/PG, 7:3 (w/w), using the Wilhelmy balance showed a rapid spreading with a surface tension of 28 mN/m after 3 s (Fig. 2). The spreading was somewhat slower using 1 weight % SP-C(LKS) in the same lipid mixture (data not shown). Addition of 2 weight % SP-B did not significantly change spreading velocity or equilibrium surface tension (Fig. 2). No improvements were observed after incubation of the mixture for 1 h at 45°C (data not shown). Similar results were obtained with DPPC:PG:PA, 68:22:9 (w/w/w) as the lipid mixture (data not shown).

Dynamic surface tension was recorded using a pulsating bubble surfactometer (Surfactometer International, Toronto, Canada) at 37°C during 50 % cyclic compression of the bubble surface and at a frequency of 40 cycles per min. All measurements were performed for 5 min and at a lipid concentration of 10 mg/ml. The pressure gradients across the bubble wall were measured at specific time intervals were measured and used to calculate surface tensions at minimum (γ_{\min}) and maximum (γ_{\max}) bubble size.

In the pulsating bubble surfactometer 3 weight % SP-C(LKS) in DPPC:PG:PA, 68:22:9 (w/w/w,) produced a surface tension of less than 1 mN/m at minimum bubble radius (γ_{\min}) while a γ_{\min} of 9-14 mN/m was observed with 3 weight % SP-C(LKS) in DPPC:PG, 7:3 (w/w) (Table 1). The surface tension at maximum bubble radius (γ_{\max}) was about 40 mN/m in both cases. Addition of 2 weight % SP-B gave γ_{\max} -values of 31-33 mN/m and γ_{\min} of 0-2 mN/m for both lipid preparations. These values are very similar to those obtained with surfactant preparations isolated from natural sources (Robertson, B. et al. (1990) Prog. Respir. Res. **25**, 237-246). Incubation of the preparations at 45°C for 1 h had no significant effect on surface activity (Table 1). Decreasing the amount of SP-B to 0.5 weight % in 3 weight % SP-C(LKS) in DPPC:PG 7:3 (w/w) tended to increase γ_{\min} although the results did not reach statistical significance (Table 1). In contrast to SP-B, addition of 2 weight % KL₄ (Cochrane, C.G. and Revak, S.D. (1991) Science **254**, 566-568) to 3 weight % SP-C(LKS) in DPPC:PG:PA 68:22:9 (w/w) did not reduce γ_{\max} which remained relatively high at 41-42 mN/m.

Example 6

Comparison between mixtures containing dipalmitoylated and non-palmitoylated reference peptides

Surfactant preparations were prepared by adding 3% w/w SP-C(Leu) or dipalmitoylated SP-C(Leu) to each lipid mixtures, made of DPPC/PG/PA 68:22:9 w/w/w. The mixtures were evaporated under nitrogen and resuspended in 150 mmol/l NaCl at lipid concentrations of 10 mg/ml. In the samples in which a SP-B substituent was also used, 1% w/w of polymyxin B was added.

Mixtures containing dipalmitoylated SP-C(Leu), with or without polymyxin B, exhibit significant improvement especially in reducing γ_{\max} at 5 min and γ_{\min} at earlier time intervals.

Table 4: Surface properties

Surface tension of the mixtures was obtained with a pulsating bubble surfactometer. After two minutes of equilibration, the recordings were obtained at different periods, at 37° C, 50% surface compression and at a rate of 40 cycles per min.

Surfactant preparation			Surface tension (mN/m)					
SP-C(Leu)	Dipalm.	PxB	7,5 s		1 min		5 min	
SP-C(Leu)								
(weight %)	(weight %)	(weight %)	γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}
1			11	39	6,2	39	2	42
1		1	3	37	3	38	0	40
	1	-	1	34	1	35	1	36
	1	1	0	29	0	34	0	35

Example 7

In vivo determination

The effect of surfactant therapy on the mechanical properties of immature lungs was evaluated in 9 preterm newborn rabbits with a gestational age of 27 days. The animals were tracheotomized at birth and five of them received, via the tracheal cannula, twice 2.5 ml/kg of artificial surfactant containing DPPC, PG, and SP-C (LKS), with or without polymyxin B, in the proportions given above. Total phospholipid concentration of the

exogenous surfactant material was 40 mg/ml. Two animals serving as negative control received no material via the tracheal tube, and another two serving as positive control were treated with the same dose of modified natural surfactant (Curosurf, Chiesi Farmaceutici Spa, Parma, Italy), diluted to 40 mg/ml. One animal was treated with a mixture of DPPC and PG in saline (same concentrations as above) at a dose of 2.5 ml/kg. All animals were kept in body plethysmograph boxes at a temperature of 37°C and ventilated in parallel for 60 min with 100% oxygen, using a Servo Ventilator 900B (Siemens-Elema, Solna, Sweden) set at a frequency of 40 min and 50% inspiration time. Tidal volumes were measured with a pneumotachygraph connected to each plethysmograph box. The animals were ventilated with a standardized tidal volume of 8-10 ml/kg and without a positive end-expiratory pressure (PEEP). Lung-thorax compliance was defined as the ratio between tidal volume and peak inspiratory pressure, and expressed as ml/cm H₂O kg.

In comparison with the non-treated control animal, compliance was improved significantly in animals treated with the artificial surfactant, especially in the animal receiving surfactant containing polymyxin B. Notably, the improvement appears to be superior to that seen after treatment with a similar dose of modified natural surfactant (Fig. 3).

Brief description of figures.

Fig. 1. Amino acid sequences and helical wheel presentations of SP-C and its analogues.

The sequence of human SP-C is taken from Johansson, J. et al. (1988) FEBS Lett. **232**, 61-64 and that of SP-C(Leu) from

Nilsson, G., et al. (1998) Eur. J. Biochem. 255, 116-124). SP-C(LKS) is based on the primary structure of SP-C but all Val residues at the positions 16-28 with the exception of position 17 are replaced with Leu residues, Lys residues have been introduced at positions 17,22 and 27, and the palmitoylated Cys at positions 5 and 6 are replaced with Ser.

Fig. 2. Surface spreading of synthetic surfactant preparations.

Spreading kinetics of 3 weight % SP-C(LKS) (filled squares, solid line) and of 3 weight % SP-C(LKS) with addition of 2 weight % SP-B (open triangles, dotted line). All preparations were examined at a concentration of 10 mg /ml of DPPC/PG, 7:3 (w/w) in 150 mmol/l NaCl. The recordings were obtained with a Wilhelmy balance and each datapoint is the mean of three different recordings.

Fig. 3 In vivo results

Lung-thorax compliance in 5 premature newborn rabbits (gestational age of 27 days) ventilated with a standardized tidal volume of 8-10 ml/kg and without a positive end-expiratory pressure (PEEP). Compliance is improved significantly in treated animals. Addition of polymyxin B (PxB) appears to increase the effect of the artificial surfactant. The concentration of phospholipids is the same in all surfactant preparations, i.e. 40 mg/ml.

09/926009

518 Rec'd PCT/PTO 13 AUG 2001

23

CLAIMS

1. SP-C analogues having general formula (I), according to one-letter amino acid code:

5 $F_e G_f I P Z Z P V H L K R (X_a B) (X_b B)_n (X_c B)_m X_d G A L L M G L$ (I)

wherein:

X is an amino acid selected from the group consisting of I, L, Nle (norleucine);

10 B is an amino acid selected from the group consisting of K, W, F, Y, Ornithine;

Z is S and can be optionally linked via ester or thio-ester bonds with acyl group containing 12-22 carbon atoms.

a is an integer from 1 to 19;

b is an integer from 1 to 19;

15 c is an integer from 1 to 21;

d is an integer from 0 to 20;

e is 0 or 1;

f is 0 or 1;

n is 0 or 1;

20 m is 0 or 1,

with the following conditions:

- $n + m > 0$;

- $f \geq e$;

25 $(X_a B) (X_b B)_n (X_c B)_m X_d$ is a sequence having a maximum of 22 amino acids, preferably from 10 to 22 amino acids.

2. SP-C analogues according to claim 1, having formula (Ia):

(Ia) FGIPSSPVHLKRX₄BX₄BX₄BXGALLMGL

3. SP-C analogues according to claim 1, having formula (Ib):

(Ib) FGIPSSPVHLKRX₅BX₅BX₄GALLMGL

4. SP-C analogues according to claim 1, having formula (Ic)

(Ic) FGIPSSPVHLKRX₄BX₁₁GALLMGL

5. SP-C analogues according to claim 1, having formula (Id)

5 (Id) FGIPSSPVHLKRX₈BX₇GALLMGL

6. SP-C analogues according to claim 1, having formula (Ie)

(Ie) FGIPSSPVHLKRX₁₁BX₄GALLMGL

7. SP-C analogues according to claims 1-6, in which Ser residues are acylated preferably with palmitoyl groups.

10 8. SP-C analogues according to claims 1-7, in which B is Lysine or Phenylalanine and X is Leucine, Isoleucine or Norleucine.

9. SP-C analogues according to claim 8, selected from the group consisting of:

SP-C (LKS) FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL

15 SP-C (LKS)₁ FGIPSSPVHLKRLILKLLLLIKLLILGALLMGL

SP-C (LKS)₂ FGIPSSPVHLKRLILKLLLLILLILGALLMGL

SP-C (LKS)₃ FGIPSSPVHLKRLILKLLLLKILLILGALLMGL

SP-C (LKS)₄ FGIPSSPVHLKRLILKLLLLIKLLILGALLMGL

SP-C (LFS) FGIPSSPVHLKRLILFLLLLFILLFLGALLMGL

20 10. A synthetic surfactant comprising at least one SP-C analogue of formula (I) in admixture with lipids and phospholipids.

11. A synthetic surfactant according to claim 10, in which the mixture lipids/phospholipids comprises DPPG, PG, PA.

12. A synthetic surfactant according to claims 10-11, further comprising
25 SP-B or an active derivative thereof or a polymyxin.

13. A synthetic surfactant according to claims 10-12, in form of solution, dispersion, suspension, dry powder.

14. Use of SP-C analogues of claims 1-7 for the preparation of a synthetic

25

surfactant to be used in all cases of surfactant deficiencies.

15. Use of a polymyxin, preferably polymyxin B for the preparation of an artificial surfactant according to claims 10-13, for the treatment of all cases of surfactant deficiencies or dysfunction, or of serous otitis media (glue ear).
- 5 16. Use according to claims 14 and 15, in which the surfactant deficiency is respiratory distress syndrome.

6092260

FIGURE 1

2/3

FIGURE 2





FIGURE 3



Declaration, Power Of Attorney and Petition

Page 1 of 3

WE (I) the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Artificial peptides having surface activity and the use thereof
in the preparation of artificial surfactant

the specification of which

☐ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and amended on _____.

☒ was filed as PCT international application

Number PCT/EP00/01044

on 09.02.2000,

and was amended under PCT Article 19

on _____ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
<u>MI99A000275</u>	<u>Italy</u>	<u>12.02.1999</u>	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No

We (I) hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

_____ (Application Number)	_____ (Filing Date)
_____ (Application Number)	_____ (Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or under § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____

And we (I) hereby appoint the following registered practitioner(s):



022850

as our (my) attorneys, with full powers of substitution and revocation, to prosecute this application and to transact all business in the Patent Office connected therewith; and we (I) hereby request that all correspondence regarding this application be sent to



022850

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1-00
CURSTEDT Tore
NAME OF FIRST SOLE INVENTOR

Tore Curstedt
Signature of Inventor


Residence: Via Palermo, 26/A

PARMA, Italy *FIX*


Citizen of: Sweden

Mailing Address: same as above

31.08.2001
Date



2-00
JOHANSSON Jan
NAME OF SECOND JOINT INVENTOR


Signature of Inventor

31.08.2001


Date

Residence: Via Palermo, 26/A
PARMA, Italy ITX

Citizen of: Sweden

Post Office Address: same as above

3-00
JÖRNVALL Hans
NAME OF THIRD JOINT INVENTOR


Signature of Inventor

31.08.2001


Date

Residence: Via Palermo, 26/A
PARMA, Italy ITX

Citizen of: Sweden

Post Office Address: same as above

3-00
ROBERTSON Bengt
NAME OF FOURTH JOINT INVENTOR


Signature of Inventor

31.08.2001


Date

Residence: Via Palermo, 26/A
PARMA, Italy ITX

Citizen of: Sweden

Post Office Address: same as above

4-00
VENTURA Paolo
NAME OF FIFTH JOINT INVENTOR


Signature of Inventor

31.08.2001

Date

Residence: Via Palermo, 26/A
PARMA, Italy ITX

Citizen of: Italy

Post Office Address: same as above

SEQUENCE LISTING

<110> Chiesi Farmaceutici spa

<120> Artificial peptides having surface activity and the use thereof in the preparation of artificial surfactant

<130> chiesi

<140>

<141>

<150> MI99A000275

<151> 1999-02-12

<160> 6

<170> PatentIn Ver. 2.1

<210> 1

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic peptides

<400> 1

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Lys Leu Leu Leu Leu Lys Ile Leu Leu Leu Lys Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 2

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic peptides

<400> 2

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Lys Leu Leu Leu Leu Ile Lys Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 3

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 3

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Lys Leu Leu Leu Leu Leu Ile Leu Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 4

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 4

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Leu Leu Leu Lys Leu Ile Leu Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu

00926000 1210

35

<210> 5

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 5

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15Leu Leu Leu Leu Leu Ile Lys Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30Met Gly Leu
35

<210> 6

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 6

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15Phe Leu Leu Leu Leu Phe Ile Leu Leu Leu Phe Leu Gly Ala Leu Leu
20 25 30Met Gly Leu
35

SEQUENCE LISTING

<110> Curstedt, Tore
Johansson, Jan
Jornvall, Hans
Robertson, Bengt
Ventura, Paolo

<120> Artificial peptides having surface activity and the use thereof in the preparation of artificial surfactant

<130> 211596US-3591-4158-0-PCT

<140> PCT/US 09/926009
<141> 2001-08-13

<150> IT MI99A000275
<151> 1999-02-12

<160> 14

<170> PatentIn version 3.1

<210> 1
<211> 101
<212> PRT
<213> ARTIFICIAL SEQUENCE

<220>
<223> SYNTHETIC PEPTIDE

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> X = Phe and may or may not be present

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> X = Gly and may or may not be present

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> X = Ser, Cys, Phe where Ser and Cys are optionally linked via ester or thio-ester bonds with acyl groups containing 12 to 22 carbons atoms linked

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> X = Ser, Cys, Phe where Ser and Cys are optionally linked via ester or thio-ester bonds with acyl groups containing 12 to 22 carbons atoms linked

09/926009 PCT/PTO

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (21)..(21)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>

<221> MISC_FEATURE
<222> (22)..(22)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (23)..(23)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (25)..(25)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (26)..(26)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (27)..(27)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (28)..(28)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (29)..(29)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (30)..(30)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (31)..(31)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (32)..(32)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (34)..(34)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (35)..(35)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (36)..(36)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (37)..(37)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (38)..(38)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (39)..(39)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (40)..(40)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (41)..(41)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>

<221> MISC_FEATURE
<222> (42)..(42)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (43)..(43)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (44)..(44)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (45)..(45)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (46)..(46)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (47)..(47)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (48)..(48)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (49)..(49)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (50)..(50)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (51)..(51)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (52)..(52)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (54)..(54)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (55)..(55)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (56)..(56)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (57)..(57)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (58)..(58)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (59)..(59)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (60)..(60)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (61)..(61)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>

<221> MISC_FEATURE
<222> (62)..(62)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (63)..(63)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (64)..(64)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (65)..(65)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (66)..(66)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (67)..(67)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (68)..(68)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (69)..(69)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (70)..(70)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (71)..(71)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (72)..(72)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (73)..(73)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (74)..(74)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (76)..(76)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (77)..(77)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (78)..(78)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (79)..(79)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (80)..(80)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (81)..(81)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>

<221> MISC_FEATURE
<222> (82)..(82)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (83)..(83)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (84)..(84)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (85)..(85)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (86)..(86)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (87)..(87)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (88)..(88)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (89)..(89)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (90)..(90)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (91)..(91)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (92)..(92)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (93)..(93)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (94)..(94)
<223> X = Val, Leu, Cys, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (95)..(95)
<223> X = Val, Leu, Ile or Nle and may or may not be present

<220>
<221> MISC_FEATURE
<222> (33)..(33)
<223> X = Ornithine, Lys, Ile, Trp, Phe, Tyr, Gln, or Asn and may or may not be present

<220>
<221> MISC_FEATURE
<222> (53)..(53)
<223> X = Ornithine, Lys, Ile, Trp, Phe, Tyr, Gln, or Asn and may or may not be present

<220>
<221> MISC_FEATURE
<222> (75)..(75)
<223> X = Ornithine, Lys, Ile, Trp, Phe, Tyr, Gln, or Asn and may or may not be present

<220>
<221> MISC_FEATURE
<222> (75)..(75)
<223> X = Ornithine, Lys, Ile, Trp, Phe, Tyr, Gln, or Asn and may or may not be present

<400> 1

Xaa Xaa Ile Pro Xaa Xaa Pro Val His Leu Lys Arg Xaa Xaa Xaa Xaa

1

5

10

15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 20 25 30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 35 40 45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 50 55 60

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 65 70 75 80

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Gly Ala
 85 90 95

Leu Leu Met Gly Leu
 100

<210> 2
 <211> 35
 <212> PRT
 <213> ARTIFICIAL SEQUENCE

<220>
 <223> SYNTHETIC PEPTIDE

<220>
 <221> MISC_FEATURE
 <222> (13)..(13)
 <223> X = Lys or Phe

<220>
 <221> MISC_FEATURE
 <222> (18)..(18)
 <223> X = Lys or Phe

<220>
 <221> MISC_FEATURE
 <222> (23)..(23)
 <223> X = Lys or Phe

<220>
 <221> MISC_FEATURE
 <222> (28)..(28)
 <223> X = Lys or Phe

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (21)..(21)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (22)..(22)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> X = Ley, Ile, or Nle

<220>

<221> MISC_FEATURE
<222> (25)..(25)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (26)..(26)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (27)..(27)
<223> X = Ley, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (29)..(29)
<223> X = Ley, Ile, or Nle

<400> 2

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Xaa Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 3
<211> 35
<212> PRT
<213> ARTIFICIAL SEQUENCE

<220>
<223> SYNTHETIC PEPTIDE

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> X = Lys or Phe

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> X = Lys or Phe

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (21)..(21)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (22)..(22)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (23)..(23)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (25)..(25)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (26)..(26)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (27)..(27)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (28)..(28)

<223> X = Leu, Ile, or Nle

<400> 3

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Xaa Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 4

<211> 35

<212> PRT

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC PEPTIDE

<220>

<221> MISC_FEATURE

<222> (17)..(17)

<223> X = Lys or Phe

<220>

<221> MISC_FEATURE

<222> (13)..(13)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (21)..(21)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (22)..(22)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (23)..(23)
<223> X = Leu, Ile, or Nle

<220>
 <221> MISC_FEATURE
 <222> (24)..(24)
 <223> X = Leu, Ile, or Nle

<220>
 <221> MISC_FEATURE
 <222> (25)..(25)
 <223> X = Leu, Ile, or Nle

<220>
 <221> MISC_FEATURE
 <222> (26)..(26)
 <223> X = Leu, Ile, or Nle

<220>
 <221> MISC_FEATURE
 <222> (27)..(27)
 <223> X = Leu, Ile, or Nle

<220>
 <221> MISC_FEATURE
 <222> (28)..(28)
 <223> X = Leu, Ile, or Nle

<400> 4
 Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Xaa Xaa Xaa Xaa
 1 5 10 15
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Gly Ala Leu Leu
 20 25 30
 Met Gly Leu
 35

<210> 5
 <211> 35
 <212> PRT
 <213> ARTIFICIAL SEQUENCE

<220>
 <223> SYNTHETIC PEPTIDE

<220>
 <221> MISC_FEATURE
 <222> (21)..(21)
 <223> X = Lys or Phe

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> X = Lue, Ile, Nle

<220>
<221> MISC_FEATURE
<222> (22)..(22)
<223> X = Lue, Ile, Nle

<220>

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> X = Lys or Phe

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> X = Leu, Ile, or Nle

<220>
<221> MISC_FEATURE
<222> (21)..(21)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (22)..(22)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (23)..(23)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (25)..(25)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (26)..(26)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (27)..(27)

<223> X = Leu, Ile, or Nle

<220>

<221> MISC_FEATURE

<222> (28)..(28)

<223> X = Leu, Ile, or Nle

<400> 6

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Xaa Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 7

<211> 35

<212> PRT

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC PEPTIDE

<400> 7

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Lys Leu Leu Leu Leu Lys Ile Leu Leu Lys Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 8

<211> 35

<212> PRT

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC PEPTIDE

<400> 8

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Lys Leu Leu Leu Leu Ile Lys Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 9

<211> 35

<212> PRT

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC PEPTIDE

<400> 9

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Lys Leu Leu Leu Leu Leu Ile Leu Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 10
<211> 35
<212> PRT
<213> ARTIFICIAL SEQUENCE

<220>
<223> SYNTHETIC PEPTIDE

<400> 10

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Leu Leu Leu Lys Leu Ile Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 11
<211> 35
<212> PRT
<213> ARTIFICIAL SEQUENCE

<220>
<223> SYNTHETIC PEPTIDE

<400> 11

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Leu Leu Leu Leu Leu Ile Lys Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 12
<211> 35
<212> PRT
<213> ARTIFICIAL SEQUENCE

<220>
<223> SYNTHETIC PEPTIDE

00999009 3440
T01111 60092550

<400> 12

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Phe Leu Leu Leu Leu Phe Ile Leu Leu Leu Phe Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 13

<211> 35

<212> PRT

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC PEPTIDE

<400> 13

Phe Gly Ile Pro Cys Cys Pro Val His Leu Lys Arg Leu Leu Ile Val
1 5 10 15

Val Val Val Val Val Leu Ile Val Val Val Ile Val Gly Gln Leu Leu
20 25 30

Met Gly Leu
35

<210> 14

<211> 34

<212> PRT

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC PEPTIDE

<400> 14

Phe Gly Ile Pro Ser Ser Pro Val Leu Lys Arg Leu Leu Ile Leu Leu
1 5 10 15

Leu Leu Leu Leu Leu Ile Leu Leu Leu Ile Leu Gly Ala Leu Leu Met
20 25 30

Gly Leu